# RHEUMATOID ARTHRITIS: LOW-DOSE METHOTREXATE

Rheumatoid arthritis (RA) is a long-term condition where the body's immune system targets the lining of the joints, causing swelling and damage. This action plan supports and informs people taking methotrexate for RA and other rheumatological conditions.

#### Understand treatment aims

Early treatment with disease modifiers reduces damage in the joints and improves symptoms such as joint pain, stiffness and swelling. Choosing the right treatment can be complex and a combination of medicines may be needed.<sup>2</sup>

#### Focus on facts

**Fact** 

Myth

Myths about methotrexate can be barriers to treatment.<sup>6</sup> Knowing the facts helps people stick to their treatment and improves results.

Methotrexate takes time to work - a response to therapy can take

from getting worse.<sup>4,5</sup>

Methotrexate works straight away.

#### **Fact**

6-12 weeks.

#### Myth

# **Fact**

Methotrexate can be safely taken with non-steroidal antiinflammatory drugs (NSAIDs).

Methotrexate has been used

its introduction in the 1950s.

Methotrexate is toxic.

safely and effectively for RA since

#### Mvth

Methotrexate should not be taken alongside NSAIDs.

### Fact

People taking methotrexate for RA can safely make physical contact with pregnant women.

#### Mvth

People taking methotrexate cannot be near pregnant women.

#### **Fact**

Methotrexate is the recommended

disease-modifying medicine in RA<sup>2,3</sup>

It is usually taken as a tablet but it can be injected. <sup>2</sup>

(within 3 months of symptoms appearing) can lessen

the signs and symptoms of RA and stop the disease

Starting low-dose methotrexate treatment early

Methotrexate in RA is used at low doses – it's not considered chemotherapy at these doses.

#### Myth

Low-dose methotrexate is chemotherapy.

## Fact

Methotrexate injections can be safely self-administered.

### Myth

Self-administration of methotrexate injections is unsafe.

## Support ongoing care

#### **Blood tests**

Check treatment is working and monitor for side effects. Measure kidney and liver function and do a full blood count.<sup>2,4</sup> Over time, these tests are needed less often.

#### Clinical review

Review joint pain and physical function regularly. How often depends on how active the disease is.<sup>2</sup>

#### **Vaccinations**

Ensure pneumococcal and influenza immunisations are up to date.2

#### Other conditions

Monitor for heart disease, osteoporosis and skin cancer, as the risk of these increases with RA. Annual skin checks are recommended.2

#### Discuss lifestyle management

- ▶ Exercise and diet Treatment may not work as well for overweight people. Work towards a healthy body weight and regular exercise.
- ▶ **Smoking** Smoking can make RA more active and treatment less effective.<sup>2</sup> Stopping smoking is recommended.
- ▶ Managing at home Tasks around the home can be difficult. Arthritis Australia's booklet **At home with arthritis:** simple steps to managing in the home may be useful (arthritisaustralia.com.au/index.php/arthritis-information/ booklets.html).
- ▶ Reproductive health Adults should seek specialist advice if they plan to have children. Women should use birth control while taking methotrexate, stop methotrexate 3 months before planning a pregnancy, and avoid breastfeeding while on methotrexate.<sup>2,8</sup>





# TAKING LOW-DOSE METHOTREXATE

Share this action plan with your healthcare team to help you achieve your treatment goals.

when I take my methotrexate	when to contact my doctor
I take my methotrexate once a week on:  Day  Dose mg  For further information see Patient Information on Methotrexate (rheumatology.org.au/MethotrexateInformation) and Self Injecting Low Dose Methotrexate (rheumatology.org.au/Self-InjectingMethotrexate) on the Australian Rheumatology Association website.  When I take my folic acid	<ul> <li>Urgently         If I develop any new infections. Signs of infection include a fever, redness or painful skin or wounds.         If I develop breathing difficulties and/or a dry cough.     </li> <li>As soon as possible</li> <li>If I experience a flare-up, I will start my flare plan.</li> <li>My flare plan:</li> </ul>
when i take my fonc acid	
I must take my folic acid <u>on different days</u>	
of the week from my methotrexate.	
Day(s)	<ul><li>Regularly</li></ul>
Dosemg	To make appointments for routine tests to monitor my disease and medicines.
Taking folic acid can help reduce the side effects of methotrexate.	To check that I am up to date with my vaccines and seek
Other medicines I take for RA	advice for travel vaccines.
······	If I am taking or plan to take any other medicines, including
Other DMARDs*	over-the-counter, herbal and naturopathic medicines.
Steroids (eg, prednisone)	
Anti-inflammatory drugs	Further information
Pain relievers	Arthritis Australia website (empowered.org.au/)
Other	Australian Rheumatology Association website
Notes/advice	( <u>rheumatology.org.au/patients/medication-information.asp</u> )
	Watch a video on how to inject methotrexate safely
* disease-modifying antirheumatic drugs	(vimeo.com/vichealth/ara)
Side effects of methotrexate	Visit the NPS MedicineWise website (nps.org.au/managing-ra)
Like all medicines, methotrexate may cause side effects.	Download the MedicineWise app to keep track of your
Most common side effects include:  ▶ nausea, vomiting, diarrhoea	medicines and access health information such as blood test results. (medicinewiseapp.com.au)

© 2018 NPS MedicineWise. Published March 2018. ABN 61 082 034 393 Level 7/418A Elizabeth St, Surry Hills NSW 2010. Independent. Not-for-profit. Evidence-based. Developed with funding from the Australian Government Department of Health. The information provided is not medical advice. Do not use it to treat or diagnose your own or another person's medical condition and never ignore medical advice or delay seeking it because of something herein. Medicines information changes, and may not be accurate when you access it. To the fullest extent permitted by law, NPS MedicineWise disclaims all liability (including without limitation for negligence) for any loss, damage, or injury resulting from reliance on, or use of this information. Any references to brands should not be taken as an endorsement by NPS MedicineWise. NPS1938.

Call the NPS Medicines Line: 1300 633 424

Date of creation:

Next review due:\_



▶ mouth ulcers

increased skin sensitivity to the sun.tiredness, headache and feeling foggy

Talk to your doctor if you are concerned. Side effects may be

reduced by taking methotrexate with food or in the evening.



